

SOME RESULTS IN GENERALIZED ŠERSTNEV SPACES

M. ALIMOHAMMADY AND R. SAADATI

ABSTRACT. In this paper, we show that D-compactness in Generalized Šerstnev spaces implies D-boundedness and as in the classical case, a D-bounded and closed subset of a characteristic Generalized Šerstnev is not D-compact in general. Finally, in the finite dimensional Generalized Šerstnev spaces a subset is D-compact if and only if it is D-bounded and closed.

1. INTRODUCTION AND PRELIMINARIES

Probabilistic normed spaces (PN spaces henceforth) were introduced by Šerstnev in 1963 [8]. In the sequel, we adopt the new definition of Generalized Šerstnev PN spaces given in the paper by Lafuerza-Guillén and Rodríguez [6]. The notations and concepts used are those of [2, 3, 4, 6] and [9].

In the sequel, the space of probability distribution functions is denoted by $\Delta^+ = \{F : \mathbb{R} \rightarrow [0, 1] : F \text{ is left-continuous, non-decreasing, } F(0) = 0 \text{ and } F(+\infty) = 1\}$ and $D^+ \subseteq \Delta^+$ is defined as follows

$$D^+ = \{F \in \Delta^+ : l^-F(+\infty) = 1\}.$$

The space Δ^+ is partially ordered by the usual point-wise ordering of functions i.e., $F \leq G$ if and only if $F(x) \leq G(x)$ for all x in $\mathbb{R}$. The maximal element for Δ^+ in this order is ε_0 , a distributive defined by

$$\varepsilon_0 = \begin{cases} 0 & \text{if } x \leq 0 \\ 1 & \text{if } x > 0. \end{cases}$$

A triangle function is a binary operation on Δ^+ , namely a function $\tau : \Delta^+ \times \Delta^+ \rightarrow \Delta^+$ that is associative, commutative, nondecreasing and which has ε_0 as unit, viz. For all $F, G, H \in \Delta^+$, we have

$$\begin{aligned} \tau(\tau(F, G), H) &= \tau(F, \tau(G, H)), \\ \tau(F, G) &= \tau(G, F), \\ F \leq G &\implies \tau(F, H) \leq \tau(G, H), \\ \tau(F, \varepsilon_0) &= F. \end{aligned}$$

Continuity of a triangle functions means continuity with respect to the topology of weak convergence in Δ^+ .

Typical continuous triangle functions are $\tau_T(F, G)(x) = \sup_{s+t=x} T(F(s), G(t))$ and $\tau_{T^*}(F, G) = \inf_{s+t=x} T^*(F(s), G(t))$.

Here, T is a continuous t-norm, i.e. a continuous binary operation on $[0, 1]$ that is commutative, associative, nondecreasing in each variable and has 1 as identity,

2000 *Mathematics Subject Classification.* 54E70.

Key words and phrases. Probabilistic normed space, Generalized Šerstnev spaces, finite dimensional space, compactness.

and T^* is a continuous t-conorm, namely a continuous binary operation on $[0, 1]$ which is related to the continuous t-norm T through $T^*(x, y) = 1 - T(1 - x, 1 - y)$.

Definition 1.1. A probabilistic normed (briefly PN) space is a quadruple (V, ν, τ, τ^*) , where V is a real vector space, τ and τ^* are continuous triangle functions, and ν is a mapping from V into Δ^+ such that, for all p, q in V , the following conditions hold:

- (N1) $\nu_p = \varepsilon_0$ if and only if, $p = \theta$, where θ is the null vector in V ;
- (N2) $\nu_{-p} = \nu_p$, for each $p \in V$;
- (N3) $\nu_{p+q} \geq \tau(\nu_p, \nu_q)$, for all $p, q \in V$;
- (N4) $\nu_p \leq \tau^*(\nu_{\alpha p}, \nu_{(1-\alpha)p})$, for all α in $[0, 1]$.

If the inequality (N4) is replaced by the equality

$$\nu_p = \tau_M(\nu_{\alpha p}, \nu_{(1-\alpha)p}),$$

then the PN space is called *Šerstnev* space and, as a consequence, a condition stronger than (N2) holds, namely

$$\nu_{\lambda p}(x) = \nu_p\left(\frac{x}{|\lambda|}\right),$$

for all $p \in V$, $\lambda \neq 0$ and $x \in \mathbb{R}$.

Following [1, 6], for $0 < b \leq +\infty$, let M_b be the set of m -transforms consist on all continuous and strictly increasing functions from $[0, b]$ onto $[0, +\infty]$. More generally, let $\widetilde{M}$ be the set of non decreasing left-continuous functions $\phi : [0, +\infty] \longrightarrow [0, +\infty]$, $\phi(0) = 0$, $\phi(+\infty) = +\infty$ and $\phi(x) > 0$, for $x > 0$. Then $M_b \subseteq \widetilde{M}$ once m is extended to $[0, +\infty]$ by $m(x) = +\infty$ for all $x \geq b$. Note that a function $\phi \in \widetilde{M}$ is bijective if and only if $\phi \in M_{+\infty}$. Sometimes, the probabilistic norms ν and ν' of two given PN spaces satisfy $\nu' = \nu\phi$ for some $\phi \in M_{+\infty}$, non necessarily bijective. Let $\hat{\phi}$ be the (unique) quasi-inverse of ϕ which is left-continuous. Recall from [9] page 49, that $\hat{\phi}$ is defined by $\hat{\phi}(0) = 0$, $\hat{\phi}(+\infty) = +\infty$ and $\hat{\phi}(t) = \sup\{u : \phi(u) < t\}$ for all $0 < t < +\infty$. It follows that $\hat{\phi}(\phi(x)) \leq x$ and $\phi(\hat{\phi}(y)) \leq y$ for all x and y .

Definition 1.2. [6]. A quadruple (V, ν, τ, τ^*) satisfying the ϕ -Šerstnev condition

$$\nu_{\lambda p}(x) = \nu_p\left(\hat{\phi}\left(\frac{\phi(x)}{|\lambda|}\right)\right),$$

for all $x \in \mathbb{R}^+$, $p \in V$ and $\lambda \in \mathbb{R} - \{0\}$ is called a ϕ -Šerstnev PN space (*Generalized Šerstnev spaces*).

Lemma 1.3. If $|\alpha| \leq |\beta|$, then $\nu_{\beta p} \leq \nu_{\alpha p}$ for every p in V .

Definition 1.4. Let (V, ν, τ, τ^*) be a PN space. For each p in V and $\lambda > 0$, the strong λ -neighborhood of p is the set

$$N_p(\lambda) = \{q \in V : \nu_{p-q}(\lambda) > 1 - \lambda\},$$

and the strong neighborhood system for V is the union $\bigcup_{p \in V} \mathcal{N}_p$ where $\mathcal{N}_p = \{N_p(\lambda) : \lambda > 0\}$.

The strong neighborhood system for V determines a Hausdorff topology for V .

Definition 1.5. Let (V, ν, τ, τ^*) be a PN space, a sequence $\{p_n\}_n$ in V is said to be strongly convergent to p in V if for each $\lambda > 0$, there exists a positive integer N such that $p_n \in N_p(\lambda)$, for $n \geq N$. Also the sequence $\{p_n\}_n$ in V is called strongly Cauchy sequence if for every $\lambda > 0$ there is a positive integer N such that $\nu_{p_n - p_m}(\lambda) > 1 - \lambda$, whenever $m, n > N$. A PN space (V, ν, τ, τ^*) is said to be strongly complete in the strong topology if and only if every strongly Cauchy sequence in V is strongly convergent to a point in V .

Definition 1.6. [4]. Let (V, ν, τ, τ^*) be a PN space and A be the nonempty subset of V . The probabilistic radius of A is the function R_A defined on $\mathbb{R}^+$ by

$$R_A(x) = \begin{cases} l^- \inf \{ \nu_p(x) : p \in A \} & \text{if } x \in [0, +\infty) \\ 1 & \text{if } x = +\infty. \end{cases}$$

Definition 1.7. [4]. A nonempty set A in a PN space (V, ν, τ, τ^*) is said to be:

- (a) certainly bounded, if $R_A(x_0) = 1$ for some $x_0 \in (0, +\infty)$;
- (b) perhaps bounded, if one has $R_A(x) < 1$, for every $x \in (0, +\infty)$ and $l^- R_A(+\infty) = 1$;
- (c) perhaps unbounded, if $R_A(x_0) > 0$ for some $x_0 \in (0, +\infty)$ and $l^- R_A(+\infty) \in (0, 1)$;
- (d) certainly unbounded, if $l^- R_A(+\infty) = 0$ i.e., if $R_A = \varepsilon_\infty$.

Moreover, A is said to be distributionally bounded, or simply D-bounded if either (a) or (b) holds, i.e. $R_A \in D^+$. If $R_A \in \Delta^+ - D^+$, A is called D-unbounded.

Theorem 1.8. [4]. A subset A in the PN space (V, ν, τ, τ^*) is D-bounded if and only if there exists a d.f. $G \in D^+$ such that $\nu_p \geq G$ for every $p \in A$.

Definition 1.9. [4]. A subset A of TVS V is said to be topologically bounded if for every sequence $\{\alpha_n\}$ of real numbers that converges to zero as $n \rightarrow +\infty$ and for every $\{p_n\}$ of elements of A , one has $\alpha_n p_n \rightarrow \theta$, in the strong topology.

Theorem 1.10. [3]. Suppose (V, ν, τ, τ^*) is a PN space, when it is endowed with the strong topology induced by the probabilistic norm ν . Then it is a topological vector space if and only if for every $p \in V$ the map from $\mathbb{R}$ into V defined by

$$\alpha \mapsto \alpha p$$

is continuous.

The PN space (V, ν, τ, τ^*) is called *characteristic* whenever $\nu(V) \subseteq D^+$.

Theorem 1.11. [6]. Let $\phi \in \widetilde{M}$ such that $\lim_{x \rightarrow \infty} \hat{\phi}(x) = \infty$. Then a ϕ -Šerstnev PN space (V, ν, τ, τ^*) is a TVS if and only if it is characteristic.

Lemma 1.12. [5, 6]. Let $\phi \in \widetilde{M}$ such that $\lim_{x \rightarrow \infty} \hat{\phi}(x) = \infty$. Let (V, ν, τ, τ^*) be a characteristic ϕ -Šerstnev PN space. Then for a subset A of V the following are equivalent

- (a) For every $n \in \mathbb{N}$, there is a $k \in \mathbb{N}$ such that $A \subset kN_\theta(1/n)$.
- (b) A is D-bounded.
- (c) A is topologically bounded.

2. D-BOUNDED AND D-COMPACT SETS IN ϕ -ŠERSTNEV SPACES

Theorem 2.1. *Let $\phi \in \widetilde{M}$ such that $\lim_{x \rightarrow \infty} \hat{\phi}(x) = \infty$. Then in a characteristic ϕ -Šerstnev PN space (V, ν, τ, τ^*) if $p_m \rightarrow p$ in V and $A = \{p_m : m \in \mathbb{N}\}$, then A is a D-bounded subset of V .*

Proof. Let $p_m \rightarrow p_0$ and $\alpha_m \rightarrow 0$. Then there exists $m_0 \in \mathbb{N}$ such that for every $m \geq m_0$, $0 < \alpha_m < 1$, then

$$\begin{aligned} \nu_{\alpha_m p_m} &\geq \tau(\nu_{\alpha_m(p_m - p_0)}, \nu_{\alpha_m p_0}) \\ &> \tau(\nu_{p_m - p_0}, \nu_{\alpha_m p_0}) \\ &\rightarrow \tau(\varepsilon_0, \varepsilon_0) \\ &= \varepsilon_0, \end{aligned}$$

as m tend to infinity. □

Example 2.2. *The 4-tuple $(\mathbb{R}, \nu, \tau_\pi, \tau_M)$, where $\nu : \mathbb{R} \rightarrow \Delta^+$ is defined by*

$$\nu_p(x) = \begin{cases} 0 & \text{if } x = 0 \\ \frac{ax}{x+|p|} & \text{if } 0 < x < +\infty \\ 1 & \text{if } x = +\infty \end{cases},$$

$a \in (0, 1)$, $\nu_0 = \varepsilon_0$ and $\phi(x) = x$, is a ϕ -Šerstnev space (see [6]). The sequence $\{\frac{1}{m}\}$ is convergent but $A = \{\frac{1}{m} : m \in \mathbb{N}\}$ is not D-bounded set. The only D-bounded set in this space is $\{0\}$.

Definition 2.3. The characteristic ϕ -Šerstnev PN space (V, ν, τ, τ^*) is said to be distributionally compact (simply D-compact) if every sequence $\{p_m\}_m$ in V has a convergent subsequence $\{p_{m_k}\}$. A subset A of a characteristic ϕ -Šerstnev PN space (V, ν, τ, τ^*) is said to be D-compact if every sequence $\{p_m\}$ in A has a subsequence $\{p_{m_k}\}$ convergent to a element $p \in A$.

Lemma 2.4. *A D-compact subset of a characteristic ϕ -Šerstnev PN space (V, ν, τ, τ^*) , is D-bounded and closed.*

Proof. Suppose that $A \subseteq V$ is D-compact. From Lemma 1.12 it is enough show that A is topologically bounded. On the contrary there is a sequence $\{p_m\} \subseteq A$ and a real sequence $\alpha_m \rightarrow 0$ such that $\{\alpha_m p_m\}$ doesn't tend to the origin in V . Then there is an infinite set $J \subseteq \mathbb{N}$ such that the sequence $\{\alpha_m p_m\}_{m \in J}$ lies in the complement of a neighborhood of the origin. Now $\{p_m\}$ is a subset of D-compact set A , so it has a convergent subsequence $\{p_m\}_{m \in J'}$. From Lemmas 1.12 and 2.1 $\{p_m\}_{m \in J'}$ is topologically bounded and so $\{\alpha_m p_m\}_{m \in J'}$ tends to origin which is a contradiction. The closedness of A is trivial. □

As in the classical case, a D-bounded and closed subset of a characteristic ϕ -Šerstnev is not D-compact in general, as one can see from the next example.

Example 2.5. *We consider quadruple $(\mathbb{Q}, \nu, \tau_\pi, \tau_M)$, where $\pi(x, y) = xy$, for every $x, y \in [0, 1]$ and probabilistic norm $\nu_p(t) = \frac{t}{t+|p|}$. It is straightforward to check that $(\mathbb{Q}, \nu, \tau_\pi, \tau_M)$ is a characteristic ϕ -Šerstnev PN space. In this space, convergence of a sequence is equivalent to its convergence in $\mathbb{R}$. We consider the subset $A = [a, b] \cap \mathbb{Q}$ which $a, b \in \mathbb{R} - \mathbb{Q}$. Since $R_A(t) = \frac{t}{t+\max\{|a|, |b|\}}$, then A is D-bounded set and since A is closed in $\mathbb{Q}$ classically, and so is closed in $(\mathbb{Q}, \nu, \tau_\pi, \tau_M)$. We*

know A is not classically compact in $\mathbb{Q}$, i.e. there exists a sequence in $\mathbb{Q}$ with no convergent subsequence in classical sense and so in $(\mathbb{Q}, \nu, \tau_\pi, \tau_M)$. Hence A is not D -compact.

Theorem 2.6. [7]. Consider a finite dimensional characteristic ϕ -Šerstnev PN space (V, ν, τ, τ^*) on real field $(\mathbb{R}, \nu', \tau', \tau'^*)$. Every subset A of V is D -compact if and only if A is D -bounded and closed.

REFERENCES

- [1] C. Alsina and B. Schweizer, The countable product of probabilistic metric spaces, *Houston J. Math.* **9** (1983), 303-310.
- [2] C. Alsina, B. Schweizer and A. Sklar, On the definition of a probabilistic normed space, *Aequationes Math.*, **46** (1993) 91-98.
- [3] C. Alsina, B. Schweizer and A. Sklar, Continuity properties of probabilistic norms, *J. Math. Anal. Appl.*, **208** (1997) 446-452.
- [4] B. Lafuerza-Guillén, A. Rodríguez-Lallena and C. Sempi, A study of boundedness in probabilistic normed spaces. *J. Math. Anal. Appl.*, **232** (1999) 183-196.
- [5] B. Lafuerza-Guillén, A. Rodríguez Lallena and C. Sempi, Normability of probabilistic normed spaces, to appear in *J. Math. Anal. Appl.*
- [6] B. Lafuerza-Guillén, Jose L. Rodríguez, Probabilistic normed spaces with non necessarily continuous triangle functions, <http://front.math.ucdavis.edu/math.PR/0408207>.
- [7] R. Saadati and M. Amini, D-bounded and D-compact sets in finite dimensional probabilistic normed spaces, to appear.
- [8] A. N. Šerstnev, On the motion of a random normed space, *Dokl. Akad. Nauk SSSR* **149**(2)(1963), 280-283(English translation in *Soviet Math. Dokl.* **4**(1963)388-390)
- [9] B. Schweizer and A. Sklar, Probabilistic Metric Spaces, Elsevier North Holland, New York, 1983.

DEPT. OF MATH., AZAD UNIVERSITY, NUR, IRAN
E-mail address: amohsen@umz.ac.ir

DEPT. OF MATH., AZAD UNIVERSITY, AMOL, P.O.BOX 678, IRAN
E-mail address: rsaadati@eml.cc